

The Dynamical Implications of Changes in mid-Stratospheric Ozone since 1991

Gerald E. Nedoluha, David E. Siskind, R. Michael Gomez
(Naval Research Laboratory)


Alyn Lambert (Jet Propulsion Laboratory)

Chris Boone (University of Waterloo, Waterloo, Canada)


Alan Parrish (University of Massachusetts, Amherst)

Ian S. Boyd (BC Scientific Consulting)

Published O₃ trends




Kyrola et al.,
ACP 2013




Gebhardt et al.,
ACP 2014

O₃ recovery in most regions, but
large long-term O₃ decrease near 30km in tropics

HALOE O₃ trend 1991-2005 (ppmv/yr)



MLS O₃ trend 2004-2013 (ppmv/yr)



Peak observed O₃ trends (~10 hPa):


~60 ppbv/yr for HALOE

~100 ppbv/yr for MLS

Some emission scenario calculations:

20 ppbv increase in surface N₂O over 20 year (IPCC A1B) => 5-7ppbv/yr decrease in O₃


Portmann et al., 2012



Annual average anomalies shown 4-times per year.
MLS is shifted to match up with HALOE during overlap period.
Decrease seems unrelated to solar cycle.

The *Positive* Correlation Between O₃ and N₂O

Monthly anomalies of
MLS O₃ and N₂O 5S-5N



Understanding the relationship between N_2O and O_3

ACE-FTS measurements


10S-10N 30km

O_3

N_2O

$\text{NO}_y = \text{NO} + \text{NO}_2 + \text{HNO}_3 + 2 * \text{N}_2\text{O}_5$

ACE sunrise and sunset measurements are both shown here





N_2O anti-correlates with NO_y

NO_y anti-correlates with O_3

→ N_2O correlates with O_3


Vertical profiles of tropical O_3 and N_2O changes from linear trends

MLS 5S-5N
2004-2013



HALOE timeseries 5S-5N, 10 hPa

HALOE O_3




HALOE does not provide N_2O ,
but it does provide NO and NO_2

O_3 and $NO+NO_2$ annual average
anomalies shown 4-times per year.


Sunset —
Sunrise - - - - -

HALOE $NO+NO_2$



$NO+NO_2$ vs. O_3 shows a
strong anti-correlation
over long timescales.

Vertical profiles of tropical O_3 and NO_x changes from linear trends



Simulating changes in tropical upwelling.


Need to compare model with fast upwelling and slow upwelling

Use simple 2D model (Bacmeister et al., 1998)

- 1) Baseline model (solid)
- 2) Increased upwelling model (Added a 0.3K/day heat source at 18km at equator, dashed).


Changes at 10 hPa are similar to observed trends

Slower upwelling gives
Decreased N_2O
Increased NO_y
Decreased O_3




What about outside the tropics?


Lauder (45S) Ozone



NDACC Microwave Ozone
Profiling Instrument
(MOPI1)
Lidar
Sonde




Unusually high 10 hPa O_3 from 2009-2013.
Large increase in mid-2009. Large decrease in mid-2013.
Good agreement between MOPI1 and MLS.




MOPI1
MLS

N_2O outside the tropics


N_2O - O_3 monthly anomaly correlation



MLS N_2O trend ppbv/yr 2004-2013



MLS O_3 trend 2004-2013 ppmv/yr



N_2O and O_3 trends from MLS since 2004 suggest:

- Older air at 10 hPa at equator
- Younger air in Southern Hemisphere between 20 hPa and 10 hPa

No such signature of younger air in HALOE O_3 (1991-2005)

MLS and NDACC Ground-Based Microwave O₃ 2004-2013

MLS O₃


MOPI (Microwave O₃ Profiling Instruments) at NDACC Lauder and Mauna Loa sites

MLS N₂O

N₂O and O₃, especially in the tropics, show a strong correlation at ~10 hPa.


Lauder

50S-40S 14.67hPa




Mauna Loa


14.5N-24.5N 10.00hPa



5S-5N 10.00hPa



Prolonged period of increased
N₂O and O₃ over Lauder



MLS and NDACC Ground-Based Microwave O₃ 2004-2014

MLS O₃


MOPI (Microwave O₃ Profiling Instruments) at NDACC Lauder and Mauna Loa sites

MLS N₂O

N₂O and O₃, especially in the tropics, show a strong correlation at ~10 hPa.


Lauder

50S-40S 14.67hPa




Mauna Loa

14.5N-24.5N 10.00hPa



5S-5N 10.00hPa



Summary

- Previously reported tropical ozone decreases at 10 hPa shown to extend over 20 years (1991-2013).
- Decrease in O_3 coincides with large decrease in N_2O (MLS) and large increase in NO_x (anticorrelates with N_2O).
- Likeliest explanation is dynamical. Slower upwelling leads to
 - Decreased N_2O
 - Increased NO_x
 - Decreased O_3
- Other, interesting results:
 - 1) From 2009-2013 10 hPa O_3 over Lauder was the highest ever observed.
 - 2) Over the last decade SH mid-latitude N_2O and O_3 show opposite behavior from tropics
 - 3) In mid-2013 there was a large increase in tropical 10 hPa O_3 and a decrease at SH mid-latitudes